

Viscosity, quark gluon plasma, and string theory

D. T. Son


Institute for Nuclear Theory, University of Washington

Plan

- Viscosity
- Heavy ion collisions and the quark gluon plasma
- Gauge/gravity duality: a surprising by-product of string theory
- Viscosity through gauge/gravity duality

Viscosity


Viscosity: introduced by Claude Navier in 1822 into what would be later called the Navier-Stokes equation.




Friction force between two plates:

$$F = \eta A \frac{\partial u_x}{\partial y}$$

Viscosity and kinetic theory of gas




 Maxwell: in the kinetic theory of gases, viscosity arises from collisions between gas molecules.


Maxwell's estimate of the viscosity:

$$\eta \sim \rho v l = \text{mass density} \times \text{velocity} \times \text{mean free path}$$

Consequence: at fixed temperature viscosity is independent of pressure (density).
Contradicts expectation at the time: the denser the gas, the larger the viscosity

Viscosity and kinetic theory of gas



 Maxwell: in the kinetic theory of gases, viscosity arises from collisions between gas molecules.

Maxwell's estimate of the viscosity:

$$\eta \sim \rho v l = \text{mass density} \times \text{velocity} \times \text{mean free path}$$

Consequence: at fixed temperature viscosity is independent of pressure (density).
Contradicts expectation at the time: the denser the gas, the larger the viscosity

“Such a consequence of a mathematical theory is very startling, and the only experiment I have met with on the subject does not seem to confirm it” (1860)

Maxwell's own experiment

During the next few years Maxwell, with the help of his wife, designed and carried out his own experiment.

Reported result in 1865: viscosity of air is independent of pressure when the latter varies from about $1/60$ to one atmosphere.



A counter-intuitive behavior

- Imagine one can turn off the interaction between molecules of a gas
- According to Maxwell's formula: viscosity $\rightarrow \infty$ (large mean free path)
- Shouldn't one expect no dissipation, $\eta \rightarrow 0$?

Reason: two limits do not commute:

- The limit of infinitely weak interaction
- The hydrodynamic limit (lengths \gg mean free path)

Viscosity is well defined only when size of experiment \gg mean free path

Viscosity of liquids: a huge range

Viscosity of fluids is much poorer understood

Table 8.4.1. *Viscosities η for some common materials in units of centipoise (10^{-2} erg s/cm³).*

Substance	Temperature	Viscosity (cp)
Air	18°C	0.018
Water	0°C	1.8
Water	20°C	1
Water	100°C	0.28
Glycerin	20°C	1500
Mercury	20°C	1.6
<i>n</i> -Pentane	20°C	0.23
Argon	85K	0.28
He ⁴	4.2K	0.033
Superfluid He ⁴	< 2.1K	0
Glass		> 10 ¹⁵

Note that, by popular convention, the designation “glass” is applied to any disordered material once its viscosity exceeds 10¹⁵cp.

(from *Chaikin & Lubensky, Principles of Condensed Matter Physics*)

Viscosity spans many orders of magnitudes, despite the fact that the density does not change much.

Viscosity of liquid glycerin

T (C)	η (mPa s)
0	12070
10	3900
20	1410
30	612
40	284
50	142
60	81.3
70	50.6
80	31.9
100	14.8
120	7.8
140	4.7
167	2.8

Liquid viscosity can be very large

The pitch-drop experiment, University of Queensland



Liquid viscosity can be very large

The pitch-drop experiment, University of Queensland



● 8 drops since 1930

Liquid viscosity can be very large

The pitch-drop experiment, University of Queensland



- 8 drops since 1930
- No one has witnessed a drop fall
- Viscosity $\sim 10^{11}$ times of water

Liquid viscosity can be very large

The pitch-drop experiment, University of Queensland



- 8 drops since 1930
- No one has witnessed a drop fall
- Viscosity $\sim 10^{11}$ times of water
- 2005 Ig Nobel Prize in Physics

Purcell and Weisskopf

- E. Purcell was very puzzled by the fact that liquid viscosity spans such a huge range

Purcell and Weisskopf

- E. Purcell was very puzzled by the fact that liquid viscosity spans such a huge range
- He thought about challenging Weisskopf to explain this fact

Purcell and Weisskopf

- E. Purcell was very puzzled by the fact that liquid viscosity spans such a huge range
- He thought about challenging Weisskopf to explain this fact

Life at low Reynolds number

E. M. Purcell

Lyman Laboratory, Harvard University, Cambridge, Massachusetts 02138

(Received 12 June 1976)

Editor's note: This is a reprint (slightly edited) of a paper of the same title that appeared in the book *Physics and Our World: A Symposium in Honor of Victor F. Weisskopf*, published by the American Institute of Physics (1976). The personal tone of the original talk has been preserved in the paper, which was itself a slightly edited transcript of a tape. The figures

Am. J. Phys. **45**, 3 (1977).

Purcell and Weisskopf

- E. Purcell was very puzzled by the fact that liquid viscosity spans such a huge range
- He thought about challenging Weisskopf to explain this fact

Life at low Reynolds number

E. M. Purcell

Lyman Laboratory, Harvard University, Cambridge, Massachusetts 02138

(Received 12 June 1976)

Editor's note: This is a reprint (slightly edited) of a paper of the same title that appeared in the book *Physics and Our World: A Symposium in Honor of Victor F. Weisskopf*, published by the American Institute of Physics (1976). The personal tone of the original talk has been preserved in the paper, which was itself a slightly edited transcript of a tape. The figures

Am. J. Phys. **45**, 3 (1977).

- but he anticipated that Weisskopf would say

$$\eta \sim \exp(\#)$$

Purcell and Weisskopf

- E. Purcell was very puzzled by the fact that liquid viscosity spans such a huge range
- He thought about challenging Weisskopf to explain this fact

Life at low Reynolds number

E. M. Purcell

Lyman Laboratory, Harvard University, Cambridge, Massachusetts 02138

(Received 12 June 1976)

Editor's note: This is a reprint (slightly edited) of a paper of the same title that appeared in the book *Physics and Our World: A Symposium in Honor of Victor F. Weisskopf*, published by the American Institute of Physics (1976). The personal tone of the original talk has been preserved in the paper, which was itself a slightly edited transcript of a tape. The figures

Am. J. Phys. **45**, 3 (1977).

- but he anticipated that Weisskopf would say

$$\eta \sim \exp(\#)$$

- So he added

Purcell's question

But it's more mysterious than that, Viki, because if you look at the Chemical Rubber Handbook table you will find that there is almost no liquid with viscosity much lower than that of water. The viscosities have a big range *but they stop at the same place*. I don't understand that. That's what I'm leaving for him.¹

Purcell's question

But it's more mysterious than that, Viki, because if you look at the Chemical Rubber Handbook table you will find that there is almost no liquid with viscosity much lower than that of water. The viscosities have a big range *but they stop at the same place*. I don't understand that. That's what I'm leaving for him.¹

Viscosity of liquids is unbounded from above, but seems to be bounded from below

A version of Purcell's question: By playing with the interaction strength, can one make $\eta \rightarrow 0$?


Modern context: the quark gluon plasma

Heavy ion collisions

- Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Lab;
- Heavy ion experiments at LHC
- RHIC: 200 GeV/pair of nucleons; LHC: 2.76 TeV/pair




A typical heavy ion collision



Goal: to create and study the quark gluon plasma

The quark gluon plasma

Hadrons consists of quarks (3 quarks, or 1 quark and 1 antiquark)




protons, neutrons



mesons (pions etc)

The quark gluon plasma


Hadrons consists of quarks (3 quarks, or 1 quark and 1 antiquark)




protons, neutrons



mesons (pions etc)



hadron gas




quark-gluon plasma

Quark-gluon plasma: quarks and gluons move relatively freely

Phases of QCD


One version of the QCD phase diagram:



(Ruester et al., hep-ph/0503184)

Strongly interacting QGP


- Unambiguous identification of QGP difficult
- However, RHIC has created some strongly interacting medium
 - Suppression of back-to-back jet correlations:



- “Elliptic flow”: signals that particles push each other (more later)


Can the viscosity be calculated?

- Good: fundamental theory known: Quantum Chromodynamics (QCD)
- Bad: perturbation theory does not work



0.2
↑
We are here!

Convergence of perturbation series



$$P/P_0|_{T=2T_c} = 1 - 0.2 + 0.4 + 0.2 - 0.8 + \dots$$

Kinetic coefficients: worse convergence expected

So what to do?

- Measure the viscosity experimentally, or
- Try a different (simpler theory)


The Gauge/Gravity Duality

Maldacena 1997: stack of N D3-branes in type IIB string theory can be described in two different pictures:

As a quantum field theory describing fluctuations of the branes: $\mathcal{N} = 4$ super-Yang-Mills theory


As string theory on a curved spacetime called $AdS_5 \times S^5$

$$ds^2 = \frac{r^2}{R^2} (-dt^2 + d\mathbf{x}^2) + \frac{R^2}{r^2} dr^2 + R^2 d\Omega_5^2$$




Holography

Gauge/gravity duality: most concrete realization of the idea of holography ('t Hooft, Susskind): a theory without gravity in (3+1)D is equivalent a theory with gravity in higher dimensions




(from Maldacena, Nature 2003)


Gauge/gravity duality at finite temperature



Gauge/gravity duality at finite temperature



Gauge/gravity duality at finite temperature



“Quark gluon plasma” = black hole (in anti de-Sitter space)

Gives entropy of plasma at strong coupling (Gubser, Klebanov, Peet 1996)

A theorist's way of measuring viscosity


How do we measure the viscosity of a system?

- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system

A theorist's way of measuring viscosity

How do we measure the viscosity of a system?


- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system



A theorist's way of measuring viscosity

How do we measure the viscosity of a system?


- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system



A theorist's way of measuring viscosity

How do we measure the viscosity of a system?


- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system



A theorist's way of measuring viscosity

How do we measure the viscosity of a system?


- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system



A theorist's way of measuring viscosity

How do we measure the viscosity of a system?

- Viscosity = response of the fluid under shear
- Theorist: send gravitational wave through the system



This is the essence of Kubo's formula


$$\eta = \lim_{\omega \rightarrow 0} \frac{1}{2\omega} \int dt d\mathbf{x} e^{i\omega t} \langle [T^{xy}(t, \mathbf{x}), T^{xy}(0, \mathbf{0})] \rangle$$

Viscosity on the light of duality

Consider a graviton that falls on this stack of N D3-branes

Will be absorbed by the D3 branes.

The process of absorption can be looked at from two different perspectives:




Viscosity on the light of duality

Consider a graviton that falls on this stack of N D3-branes

Will be absorbed by the D3 branes.

The process of absorption can be looked at from two different perspectives:




Viscosity on the light of duality

Consider a graviton that falls on this stack of N D3-branes

Will be absorbed by the D3 branes.

The process of absorption can be looked at from two different perspectives:





Viscosity on the light of duality

Consider a graviton that falls on this stack of N D3-branes

Will be absorbed by the D3 branes.

The process of absorption can be looked at from two different perspectives:



Absorption by D3 branes (\sim viscosity) = absorption by black hole

Viscosity/entropy density ratio

Collecting facts so far:

- Viscosity \sim absorption cross section for low energy gravitons
- Absorption cross section = area of horizon (a consequence of Einstein equation)
- Entropy: also \sim area of horizon (Bekenstein-Hawking formula)

$$\frac{\eta}{s} = \frac{1}{4\pi}$$

where η is the shear viscosity, s is the entropy per unit volume.

Universality of η/s

Restoring \hbar and c in the formula: a surprise

$$\frac{\eta}{s} = \frac{\hbar}{4\pi}$$

- No velocity of light c
- Finite viscosity at infinitely strong coupling!
- *The same value in all theories with gravity duals* Kovtun, DTS, Starinets 2003

Universality of η/s is related to properties of black hole horizons

Why does η/s have the dimensionality of \hbar

In kinetic theory

$$\eta \sim \rho v l, \quad s \sim n = \frac{\rho}{m}$$

$$\frac{\eta}{s} \sim m v l \sim \hbar \frac{\text{mean free path}}{\text{de Broglie wavelength}}$$

- In kinetic theory, mean free path cannot be \ll de Broglie wavelength.
- Uncertainty principle: η/s bounded from below by $\# \hbar$, unknown coefficient
- Theories with gravity dual reach $\hbar/4\pi$

Why does η/s have the dimensionality of \hbar

In kinetic theory

$$\eta \sim \rho v l, \quad s \sim n = \frac{\rho}{m}$$

$$\frac{\eta}{s} \sim m v l \sim \hbar \frac{\text{mean free path}}{\text{de Broglie wavelength}}$$

- In kinetic theory, mean free path cannot be \ll de Broglie wavelength.
- Uncertainty principle: η/s bounded from below by $\# \hbar$, unknown coefficient
- Theories with gravity dual reach $\hbar/4\pi$

Conjecture [Kovtun, DTS, Starinets 2003](#)


$$\frac{\eta}{s} \gtrsim \frac{\hbar}{4\pi}$$

Reminds Purcell's question to Weisskopf

No c : $\hbar/(4\pi)$ can be compared with ordinary, nonrelativistic liquids

Ordinary liquids

η/s of liquid water in unit of $\hbar/(4\pi)$, as function of temperature (K), along the saturation curve



General observation on liquids

- η/s reaches minimum near the critical point (liquid=gas) Kapusta, McLerran
- but not exactly at the critical point: η diverges there according to theory of dynamic critical phenomena
- The minimal value of η/s varies from substance to substance


Substance	$(\frac{\eta}{s})_{\min}$	Substance	$(\frac{\eta}{s})_{\min}$	Substance	$(\frac{\eta}{s})_{\min}$
H, He	8.8				
Ne	17	H ₂ O	25	CO	35
Ar	37	H ₂ S	35	CO ₂	32
Kr	57	N ₂	23	SO ₂	39
Xe	84	O ₂	28		

(η/s is measured in unit of $\hbar/(4\pi)$)

Minimum among substances is reached by the most quantum liquids: hydrogen and helium

Superfluids: normal component finite shear viscosity (Andronikashvili experiments)


determining viscosity of QGP



- Collisions with nonzero impact parameter
- Distribution of particles over momentum is not axially symmetric: characterized by “elliptic flow” parameter v_2
- explanation: pressure gradient depends on angle

- Viscosity reduces v_2
- Hydrodynamic simulations can give estimates for η

Recent numerical simulations



(from Luzum and Romatschke, arXiv:0804. LHC data seem to prefer Glauber initial conditions.)

$$\frac{\eta}{s} = 0.1 \pm 0.1(\text{th}) \pm 0.08(\text{exp})$$

Not too far way from $1/4\pi$

QGP is strongly coupled (sQGP)

High or low viscosity?

Brookhaven National Lab press release 2005: “the degree of collective interaction, rapid thermalization, and **extremely low viscosity** of the matter being formed at RHIC make this the **most nearly perfect liquid ever observed**.”

High or low viscosity?

Brookhaven National Lab press release 2005: “the degree of collective interaction, rapid thermalization, and **extremely low viscosity** of the matter being formed at RHIC make this the **most nearly perfect liquid ever observed**.”

Estimating the viscosity of the QGP:

$$\eta \sim \frac{\hbar}{4\pi} s \sim \frac{10^{-27} \text{ erg} \cdot \text{s}}{(10^{-13} \text{ cm})^3} \sim 10^{14} \text{ cp}$$

High or low viscosity?

Brookhaven National Lab press release 2005: “the degree of collective interaction, rapid thermalization, and **extremely low viscosity** of the matter being formed at RHIC make this the **most nearly perfect liquid ever observed**.”

Estimating the viscosity of the QGP:

$$\eta \sim \frac{\hbar}{4\pi} s \sim \frac{10^{-27} \text{ erg} \cdot \text{s}}{(10^{-13} \text{ cm})^3} \sim 10^{14} \text{ cp}$$

Table 8.4.1. *Viscosities η for some common materials in units of centipoise (10^{-2} erg s/cm³).*

Substance	Temperature	Viscosity (cp)
Air	18°C	0.018
Water	0°C	1.8
Water	20°C	1
Water	100°C	0.28
Glycerin	20°C	1500
Mercury	20°C	1.6
<i>n</i> -Pentane	20°C	0.23
Argon	85K	0.28
He ⁴	4.2K	0.033
Superfluid He ⁴	< 2.1K	0
Glass		> 10 ¹⁵

Note that, by popular convention, the designation “glass” is applied to any disordered material once its viscosity exceeds 10¹⁵cp.

High or low viscosity?

Brookhaven National Lab press release 2005: “the degree of collective interaction, rapid thermalization, and **extremely low viscosity** of the matter being formed at RHIC make this the **most nearly perfect liquid ever observed**.”

Estimating the viscosity of the QGP:

$$\eta \sim \frac{\hbar}{4\pi} s \sim \frac{10^{-27} \text{ erg} \cdot \text{s}}{(10^{-13} \text{ cm})^3} \sim 10^{14} \text{ cp}$$

Table 8.4.1. *Viscosities η for some common materials in units of centipoise ($10^{-2} \text{ erg s/cm}^3$).*

Substance	Temperature	Viscosity (cp)
Air	18°C	0.018
Water	0°C	1.8
Water	20°C	1
Water	100°C	0.28
Glycerin	20°C	1500
Mercury	20°C	1.6
<i>n</i> -Pentane	20°C	0.23
Argon	85K	0.28
He ⁴	4.2K	0.033
Superfluid He ⁴	< 2.1K	0
Glass		> 10 ¹⁵


Note that, by popular convention, the designation “glass” is applied to any disordered material once its viscosity exceeds 10¹⁵cp.

In absolute terms, QGP is almost as viscous as glass!

Low viscosity: in the sense of small η/s , suggested by gauge/gravity duality

Condensed matter analogy

Mott's minimal metallic conductivity hypothesis: conductivity is bounded from below by uncertainty principle



(Phillips, *Advanced Solid State Physics*)

Mott's minimal metallic conductivity is a useful concept, but not a real, absolute minimum.

Is there a viscosity bound?

● $\frac{\eta}{s} \geq \frac{\hbar}{4\pi}$ for all fluids? Kovtun, DTS, Starinets 2005

● There exist theories where

$$\frac{\eta}{s} = \frac{\hbar}{4\pi} \left(1 - \frac{c}{N}\right), \quad N \gg 1$$

Kats, Petrov; Buchel, Myers, Sinha

● No reliable way to go to small N .

● Model studies: bad things happen before η/s reaches 0 Brigante etc. PRL 2008

● Within a model: causality is violated when $\eta/s > \frac{16}{25} \frac{\hbar}{4\pi}$

● What is the real minimum of η/s ?

Further applications

Chiral separation in quark matter:

- Rotating volume of quark matter with nonzero chemical potential
- Left and right handed quarks separate along the axis of rotation
- Seen in gauge-gravity duality, and understood to be general feature of chiral relativistic fluids
- Originates from *quantum anomalies*

Conclusion

Surprising applications of string theory to real-world problems

- New perspective on old problems
 - QGP = black hole
 - viscosity = gravity absorption
 - etc
- Suggested η/s as a relevant ratio
- Suggested the value $\hbar/(4\pi)$ as particularly interesting.

Most serious challenge: connect to the correct theory of strong interactions, QCD

The unity of physics

We have also learned a lesson that physics is one single subject:

- Hydrodynamics
- Quantum field theory
- Statistical mechanics
- Gravity
- String theory

The End