


# The Pioneer Anomaly: The Data, Its Meaning, and a Future Test



Michael Martin Nieto  
Los Alamos National Laboratory  
University of California

Physics Dept. Colloquium

University of Toronto

29 Sept. 2005

# The original Pioneer Collaboration


John D. Anderson	JPL
Phillip A. Laing	Aerospace*
Eunice L. Lau	JPL
Anthony S. Liu	Astro Sci*
Michael Martin Nieto	LANL
Slava G. Turyshev	JPL

Phys. Rev. Lett. **81**, 2858-2861 (1998), gr-qc/9808081

Phys. Rev. D **65**, 082004/1-50 (2002), gr-qc/0104064

# A) THE DATA

Pioneer F (10) at the Cape




Pioneer 10: 2 March 1972





# Meanwhile ...



# Pioneer 10/11: Main Missions




# Pioneers in the galaxy



# Early Data

UNMODELED ACCELERATIONS ON PIONEER 10 AND 11  
Acceleration Directed Toward the Sun



As preparing for 1994 talk on gravity and anti-matter (see Bled Proceedings), John emailed:

By the way, the biggest systematic in our acceleration residuals is a bias of

$8 \times 10^{-13} \text{ km/s}^2$

directed toward the Sun.

*This is 8 Angstroms/s<sup>2</sup> !!*

$$a_N = 5.93 \times 10^{-6} \text{ km/s}^2, \text{ at 1 AU}$$




# THE EXTERNAL REACTIONS

- 1) “IT MUST BE A GLITCH THAT WILL GO AWAY WITH TIME. THIS CODE WORKS!”
- 2) IT DID NOT GO AWAY. “BUT WHO CARES? IT IS SMALL AND THINGS WORK WELL ENOUGH.”
- 3) THEN WE STARTED STRONGLY ASSERTING THAT THE EFFECT REALLY IS IN THE DATA.
- 4) “WELL, IT MUST BE THE CODE AFTER ALL. DON’T BOTHER US ANY MORE UNLESS YOU SHOW US IT IS NOT THE CODE.”


... MUMBLE GRUMBLE

- 5) FINALLY ANOTHER CODE was used besides ODP... CHASMP.

# From CHASMP (Aerospace)



# ODP results (JPL)



## Error Budget: A Summary of Biases and Uncertainties

Item	Description of error budget constituents	Bias $10^{-8} \text{ cm/s}^2$	Uncertainty, $10^{-8} \text{ cm/s}^2$
1	Systematics generated external to the spacecraft:		
	a) Solar radiation pressure and mass	+0.03	$\pm 0.01$
	b) Solar wind		$\pm < 10^{-5}$
	c) Solar corona		$\pm 0.02$
	d) Electro-magnetic Lorentz forces		$\pm < 10^{-4}$
	e) Influence of the Kuiper belt's gravity		$\pm 0.03$
	f) Influence of the Earth orientation		$\pm 0.001$
	g) Mechanical and phase stability of DSN antennae		$\pm < 0.001$
	h) Phase stability and clocks		$\pm < 0.001$
	i) DSN station location		$\pm < 10^{-5}$
	j) Troposphere and ionosphere		$\pm < 0.001$
2	On-board generated systematics:		
	a) Radio beam reaction force	+1.10	$\pm 0.11$
	b) RTG heat reflected off the craft	-0.55	$\pm 0.55$
	c) Differential emissivity of the RTGs		$\pm 0.85$
	d) Non-isotropic radiative cooling of the spacecraft		$\pm 0.48$
	e) Expelled Helium produced within the RTGs	+0.15	$\pm 0.16$
	f) Gas leakage		$\pm 0.56$
	g) Variation between spacecraft determinations	+0.17	$\pm 0.17$
3	Computational systematics:		
	a) Numerical stability of least-squares estimation		$\pm 0.02$
	b) Accuracy of consistency/model tests		$\pm 0.13$
	c) Mismodeling of maneuvers		$\pm 0.01$
	d) Mismodeling of the solar corona		$\pm 0.02$
	e) Annual/diurnal terms		$\pm 0.32$
Estimate of total bias/error		+0.90	$\pm 1.33$

$$a_P = a_{P(\text{exper})} + b_P \pm \sigma_P, \quad a_{P(\text{exper})} = (7.84 \pm 0.01) \times 10^{-8} \text{ cm/s}^2$$

$$a_P = (8.74 \pm 1.33) \times 10^{-8} \text{ cm/s}^2 \sim (8.7 \pm 1.3) \times 10^{-8} \text{ cm/s}^2.$$



## Large Systematics (in units of $10^{-8}$ cm/s<sup>2</sup>)

a) Radio beam	+ 1.10	+/- 0.11
b) RTG heat reflection	- 0.55	+/- 0.55
c) Differential RTG emission		+/- 0.85
d) Thermal cooling		+/- 0.48
e) Gas leaks		+/- 0.56


One can *interpret* the Doppler frequency drift as

$$a_p = (8.74 \pm 1.33) \times 10^{-8} \text{ cm/s}^2$$

# SNAP19 RTGs




# Electrical power



1987.0 \*

1998.8 \*

# B) THE DATA'S MEANING





# What do we really “know” from the big study?

- For Pioneer 10: between ~40-70.5 AU (1987.0-1998.5)

$$a_{P(\text{expt})}^{\text{Pio 10}} = (7.84 \pm 0.01) \times 10^{-8} \text{ cm/s}^2$$

- For Pioneer 11: between ~22.4-31.7 AU (1987.0-1990.8)

$$a_{P(\text{expt})}^{\text{Pio 11}} = (8.55 \pm 0.02) \times 10^{-8} \text{ cm/s}^2$$





- Analysis for both Pioneers with systematics:


$$a_p = (8.74 \pm 1.33) \times 10^{-8} \text{ cm/s}^2$$

SEEN only on these small (~250 kg) craft on hyperbolic orbits.  
NOT SEEN on large, bound, astronomical bodies.

**But REMEMBER, this is really a Doppler shift,  
that is only INTERPRETED as an acceleration.**

# Dust Density and Drag

$a_p$    
 $\square_p$   $v_p$    $A_p$    $m_p^{19}$   
 $\square_{IPD}$    $\square_p$   $\rho$   $\text{g/cm}^3$   
 $3 \times 10^{-19} \text{ g/cm}^3$



- (a) Pioneer upper bound on IPD from drag
- (b, c) Model-dependent upper bounds on IPD
- (d) Estimate of IPD
- (e) Estimate of ISD

**BOTTOM LINE: Any drag is DARK MATTER, not dust**

# KB matter and Gravity

$$\mathbf{a}_{\text{KB}}(\mathbf{r}) \quad \text{[book icon]} \quad m_p \quad \text{[clock icon]}$$




$$d^3r' \quad (-G) \quad \rho_{\text{KB}}(\mathbf{r}') \quad /$$

A total spherical  $1/r$  density yields a constant acceleration, whereas a shell does not.

Further,  $1/r$  disk with

$$\rho_{\text{KB}} = \rho_0 / r' ; \quad 10 \text{ AU} \leq r' \leq 100 \text{ AU}; \quad 1 \text{ AU} \leq z \leq -1 \text{ AU}$$

does NOT yield a constant acceleration.



**2nd BOTTOM LINE: KB matter WILL NOT DO IT**

## What do we only “suspect” or not know?



- We have no real idea how far out the anomaly goes.
- $a_p$  continues out *roughly* as a constant from about 10 AU.

**BUT:**

- Pioneer 10 shows an “effect” starting only at  $\sim 10$  AU.
- Before Saturn encounter (at 10 AU) and the transition to hyperbolic orbit, Pioneer 11 did not show the anomaly.

# Onset of the Anomaly?


UNMODELED ACCELERATIONS ON PIONEER 10 AND 11  
Acceleration Directed Toward the Sun



At Saturn Pioneer 11 reached escape velocity and anomaly had big error. Is it a drag turning on or the escape velocity? (Pio 10 escaped at Jupiter.)


# C) A FUTURE TEST

I: **The early data** from 6/78 has been retrieved and will be properly reanalyzed. Although clouded by solar radiation pressure, it will give us more information on the time-dependence and could reveal the anomaly's direction.







- 1) Towards the Sun: gravity?
- 2) Towards the Earth: time?
- 3) Along the velocity: drag or inertia?
- 4) On the spin axis: internal systematics?

# Signals of different anomaly directions



- 1) Towards the Sun: gravity?
- 2) Towards the Earth: time?
- 3) Along the velocity: drag or inertia?
- 4) On the spin axis: internal systematics?

Retrieved data contains good Saturn encounter. Also have short data arcs around earlier Jupiter encounters.





## II: Possibilities for an add-on experiment

A. New Horizons mission to Pluto

B. Jettisoned package from InterStellar Probe?




# III: A Dedicated Mission

## LESSONS LEARNED FROM THE PIONEERS

- Spin Stabilization
- Precise Doppler navigation
- RTGs (at the ends of long booms?)
- Thermal design with low asymmetry
- Well-engineered craft and mission

We want to emphasize the systematic problems that any successful mission will have to address.




# Mission Options

- **Fore/aft symmetric deep-space mission**
- **Formation mission**
- **Accelerometer**

# FORE/AFT SYMMETRIC DESIGN PROPOSES UNIQUE FEATURES


- Symmetric fore/aft thermal design, including louvers on the sides of the central bus
- Dual fore/aft antennas
- J. D, Anderson, MMN, and S. G. Turyshev,  
Mod. Phys. D **11**, 1545-1553 (2002). gr-qc/0205059
- MMN and S G. Turyshev,  
Mod. Phys. D **13**, 899-906 (2004), gr-qc/0308108
- MMN and S. G. Turyshev,  
Class. Quant. Grav **21**, 4005-4023 (2004), gr-qc/0308017

# Proposed mission concept



## How design would kill the systematics:


- a) Broadcast in both directions so radiation force cancels.
- b) Positions of RTGs and louvers, coupled with symmetric fore/aft antenna configurations and the rotation of the craft, mean heat and power are radiated axially symmetrically fore/aft, and hence have no effect.



But what if there were some imperfection  
(like stuck louvers or a degraded antenna)?

To take care of this, after one  
year rotate the craft by 180  
degrees!

(The Pioneer 10 “Earth  
Acquisition Maneuver” took  
two hours and 0.5 kg fuel.)



$$a_p = (a_{\text{forward}} + a_{\text{backward}})/2$$



With off-the-shelf technology one could obtain

$$\diamond \sim 0.06 \times 10^{-8} \text{ cm/s}^2,$$

in a few years of data taking,

**IF**

the thrusters are reliable and gas leaks can be eliminated or monitored to a high enough accuracy.




With new technology one could reach

$\diamond \sim$

$$0.01 \times 10^{-8} \text{ cm/s}^2$$



# ESA Cosmic Vision Theme: A NEW PIONEER COLLABORATION

- H. Dittus, C. Lämmerzahl, S. Theil (ZARM, University of Bremen)
  - Bernd Dachwald, Wolfgang Seboldt (German Aerospace Center)
    - W. Ertmer, E. Rasel (University of Hanover)
      - U. Johann (Astrium Space, Germany)
  - B. Kent, R. Bingham (Rutherford Appleton Laboratory)
    - O. Bertolami (University of Lisbon)
      - T. Touboul (ONERA, France)
        - P. Bouyer (Orsay, France)
      - S. Reynaud (ENS/LKB, France)
  - C. Erd, C. de Matos, A. Rathke (ESA/ESTEC, Netherlands)
  - J. D. Anderson, S. G. Turyshev (Jet Propulsion Laboratory)
    - M. M. Nieto (Los Alamos National Laboratory (LANL))



Back (XY) Smooth Shade

Smooth Shade



# Mission Summary

## Objectives

- To search for any *unmodeled* small acceleration affecting the spacecraft motion at the level of  $\sim 0.1 \times 10^{-8} \text{ cm/s}^2$  *or less*.
- Determine the physical origin of any anomaly, if found.

## Features

- A standard spacecraft bus that allows thermal louvers to be on the sides for symmetric fore/aft thermal rejection.

# Spacecraft

- Power at launch: ~200W provided by RTGs located on booms at a distance of ~3 m from the rotational axis of the spacecraft or shielded.
- Mass: s/c dry ~300 kg; propellant ~40 kg; total at launch ~500 kg.
- Dimensions at launch: diameter ~2.5 m; height: ~3.5 m or less.
- Attitude control: spin-stabilized spacecraft.
- Navigation: Doppler, range, and possibly VLBI and/or  $\Delta$ DO R.
- m W laser to Probe.

## Orbit

- Solar system escape trajectory -- possibly in the plane of ecliptic, co-moving with the solar system's direction wrt local IS medium.
- Spacecraft moving with a velocity of 5 AU or more per year, reaching 15 AU in 3 years time or less.

## Lifetime

- 7 years (nominal for velocity of 5 AU/year); 12 years (extended).

## Launcher


- Ariane 5, Proton, or any heavy vehicle, Delta IV 2425, etc.

We want to get there quick!



- As stated, a test could be either a stand alone mission or a probe of a large mission that is jettisoned after final propulsion is over.
- Such a mission would unambiguously determine the validity of the Pioneer anomaly.
- It would also advance the metrology of deep space navigation to unprecedented levels, something that will be needed in the future.
- Independent of the anomaly this would be very important.
- But if the anomaly exists, then ...






This wide-range time exposure taken from the island of La Palma in the Canary Islands reveals an incredible view of stars, nebulae, the constellation Orion, and the Milky Way. Stretching across the image from the bottom left, faint stars compose the luminous band of the Milky Way. A group of yellowish stars at the upper right is dominated by the red giant Aldebaran, where Pioneer 10 is headed. Image: A. Vannini, G. Li Causi, A. Ricciardi, and A. Garatti

HYPERFINE TRANSITION OF  
NEUTRAL HYDROGEN

SILHOUETTE OF  
SPACECRAFT

BINARY EQUIVALENT  
OF DECIMAL 8



POSITION OF SUN  
RELATIVE TO 14  
PULSARS AND THE  
CENTER OF THE GALAXY

PLANETS OF SOLAR  
SYSTEM AND BINARY  
RELATIVE DISTANCES