

PHY152 – Practice Problem Set 8 Solutions

1) Wolfson 13.55

This problem is about simple harmonic motion of a pendulum. We are asked to solve for the period, first by treating it as a simple pendulum and then as a physical pendulum.

Period for a simple pendulum is:

$$T = 2\pi \sqrt{\frac{mL^2}{mgL}} = 2\pi \sqrt{\frac{L}{g}}$$

$$T = 2\pi \sqrt{\frac{.8}{9.8}} = 1.795 \text{ s}$$

For the physical pendulum we will need to invoke Parallel-Axis Theorem to account for the rotational inertia of the ball/sphere (Table 10.2) about the pivot point:

$$I = I_{cm} + md^2 = \frac{2}{5}mR^2 + mL^2$$

Hence,

$$T = 2\pi \sqrt{\frac{\frac{2}{5}mR^2 + mL^2}{mgL}} = 2\pi \sqrt{\frac{\frac{2}{5}R^2 + L^2}{gL}}$$

$$T = 2\pi \sqrt{\frac{\frac{2}{5} * .075^2 + .8^2}{9.8 * .8}} = 1.798 \text{ s}$$

$$\% \text{ Difference} = \left| \frac{1.798 - 1.795}{1.798} \right| 100\% = .17\%$$

2) Wolfson 13.65

This problem involves determining # of oscillations for the amplitude to decay to 1/e of its initial value in a spring-damper system. The mass (0.25kg) is mounted on a spring (3.3 N/m) and the damping constant is $b = 8.4 \times 10^{-3}$ kg/s.

Examining the equation which models the damped oscillations:

$$x(t) = Ae^{-\frac{bt}{2m}} \cos(\omega t + \phi)$$

the term responsible for decay is

$$e^{-\frac{bt}{2m}}$$

Hence,

$$e^{-\frac{bt}{2m}} = e^{-1}$$

$$\frac{bt}{2m} = 1$$

$$t = \frac{2m}{b} = \frac{2(.25)}{8.4 \times 10^{-3}} = 59.5 \text{ s}$$

Angular frequency:

$$\omega = \sqrt{\frac{k}{m}} = \sqrt{\frac{3.3}{.25}} = 3.6 \frac{\text{rad}}{\text{s}}$$

Period:

$$T = \frac{2\pi}{\omega} = 1.75 \text{ s}$$

Hence,

$$\#oscillations = \frac{t}{T} = \frac{59.5}{1.75} = \mathbf{34 \text{ oscillations}}$$

3) Wolfson 13.82

This problem involves torsional oscillation of a disk with two masses attached at the end. We are given that the disk has diameter 50cm ($R=0.25\text{m}$), mass of 340g ($M=0.34\text{kg}$) and is suspended by a wire with torsional constant ($\kappa=5\text{N m/rad}$). Once the birds land on opposite sides, frequency of oscillation is ($f=2.6\text{Hz}$).

For torsional oscillation:

$$\omega = \sqrt{\frac{\kappa}{I}}$$

Total rotational inertia will consist of:

$$I = I_{\text{disk}} + I_{\text{birds}}$$
$$I = \frac{1}{2}MR^2 + 2mR^2$$

So,

$$2\pi f = \sqrt{\frac{\kappa}{\frac{1}{2}MR^2 + 2mR^2}}$$
$$m = \frac{1}{2R^2} \left\{ \kappa \left(\frac{1}{2\pi f} \right)^2 - \frac{1}{2}MR^2 \right\}$$
$$m = \frac{1}{2(.25^2)} \left\{ 5 \left(\frac{1}{2\pi(2.6)} \right)^2 - \frac{1}{2} \cdot 34(.25^2) \right\}$$
$$m = \mathbf{0.065 \text{ kg}}$$